

# INTRODUCTION TO MODULAR FORMS

## LOTHAR GÖTTSCHE

*Stefano Maggiolo\**

*SISSA, Trieste*

*March 9<sup>th</sup>, 2009–March 16<sup>th</sup>, 2009*

### CONTENTS

1	Introduction	1
2	The modular group	2
3	Examples	4
4	Zeroes of modular forms	6
5	Theta functions	9
6	Modular forms for congruence subgroups	10
7	Hecke theory	12
8	L-series	14
	References	15

### 1 INTRODUCTION

Let  $E_\Lambda$  be the elliptic curve associated to lattice  $\Lambda = \mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2$ , oriented in the sense that  $\Im\omega_1/\omega_2 > 0$ . We know that  $E_{\Lambda_1} \cong E_{\Lambda_2}$  if and only if  $\Lambda_1 = a\Lambda_2$  for some  $a \in \mathbb{C} \setminus \{0\}$ .

*Lecture 1 (2 hours)  
March 9<sup>th</sup>, 2009*

**1.1 DEFINITION.** A *modular function* is a function  $M_{1,1} \rightarrow \mathbb{C}$  where  $M_{1,1}$  is the space of elliptic curves over  $\mathbb{C}$ .

A modular function can be viewed as a function

$$F: \{\mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2\} \rightarrow \mathbb{C}$$

with the property that  $F(a\Lambda) = F(\Lambda)$  for all lattice  $\Lambda$  and  $a \in \mathbb{C} \setminus \{0\}$ .

---

\*s.maggiolo@gmail.com

## 2. THE MODULAR GROUP

---

1.2 DEFINITION. A modular form of weight  $k$ , with  $k \in \mathbb{Z}$ , is a modular function  $F$  such that

$$(1) \quad F(a\Lambda) = a^{-k}F(\Lambda).$$

Let  $\text{SL}(2, \mathbb{Z})$  the set of integral matrices with determinant 1; from now on, we will denote an element of  $\text{SL}(2, \mathbb{Z})$  with  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ . Let  $\Lambda := \mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2$  a lattice; if  $A \in \text{SL}(2, \mathbb{Z})$ ,  $A(\omega_1, \omega_2)$  is just another basis for the same lattice. Let  $\mathbb{H} := \{\tau \in \mathbb{C} \mid \Im\tau > 0\}$  be the half complex plane with positive imaginary part. The matrix group  $\text{SL}(2, \mathbb{Z})$  acts on  $\mathbb{H}$  by  $A(\tau) := \frac{a\tau+b}{c\tau+d}$ .

Now let  $F$  be a modular form of weight  $k$ ; we can associate to it a function  $f: \mathbb{H} \rightarrow \mathbb{C}$  by  $f(\tau) := F(\mathbb{Z}\tau \oplus \mathbb{Z})$ ; in this context, condition (1) ensure that

$$(2) \quad \begin{aligned} f(A(\tau)) &= F\left(\mathbb{Z}\frac{a\tau+b}{c\tau+d} \oplus \mathbb{Z}\right) = F\left(\frac{1}{c\tau+d}(\mathbb{Z}(a\tau+b) \oplus \mathbb{Z}(c\tau+d))\right) = \\ &= (c\tau+d)^k F(A(\mathbb{Z}\tau \oplus \mathbb{Z})) = (c\tau+d)^k F(\mathbb{Z}\tau \oplus \mathbb{Z}) = \\ &= (c\tau+d)^k f(\tau). \end{aligned}$$

Conversely, if we have  $f: \mathbb{H} \rightarrow \mathbb{C}$  such that  $f(A(\tau)) = (c\tau+d)^k f(\tau)$ , then we can associate to it a function  $F$  as in  $F(\mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2) = \omega_2^{-k} f(\omega_1/\omega_2)$ ; this is a modular form. In particular one obtain that these correspondences are each the inverse of the other. So we can give the following equivalent definition.

1.3 DEFINITION. A modular form of weight  $k$ , with  $k \in \mathbb{Z}$ , is a holomorphic function  $f: \mathbb{H} \rightarrow \mathbb{C}$  satisfying (2), and not growing too fast as  $\tau \rightarrow \infty$ .

The last condition will ensure later that modular forms corresponds to sections of a line bundle on  $\overline{M}_{1,1}$ . Another way to say the same thing is to define for every  $f: \mathbb{H} \rightarrow \mathbb{C}$ ,  $k \in \mathbb{Z}$ , and  $A \in \text{SL}(2, \mathbb{Z})$  the function  $f|_{k,A}: \mathbb{H} \rightarrow \mathbb{C}$  with  $f|_{k,A}(\tau) := (c\tau+d)^{-k} f(A\tau)$ ; then we request  $f = f|_{k,A}$  for every  $A$ .

Why modular forms are useful in mathematics?

1. There are very few modular forms; the space of modular forms of weight  $k$  is a vector space of finite dimension.
2. They occur naturally in many fields of mathematics and physics.

## 2 THE MODULAR GROUP

Consider the previously defined action of  $\text{SL}(2, \mathbb{Z})$  on  $\mathbb{H}$ ; since  $-I$  acts trivially, we can also say that  $\Gamma := \text{SL}(2, \mathbb{Z})/\{\pm I\}$  acts on  $\mathbb{H}$ . We define two special elements:

1.  $S := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ , such that  $S\tau = -\tau^{-1}$ ;
2.  $T := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ , such that  $T\tau = \tau + 1$ .

Moreover, we have  $S^2 = I = (ST)^3$  in  $\Gamma$ .




Figure 1: A fundamental domain of the action of  $\Gamma$  on  $\mathbb{H}$ .

Now we can find a fundamental domain for the action of  $\Gamma$ :

$$D := \{\tau \in \mathbb{H} \mid |\tau| \geq 1, -1/2 \leq \Re\tau \leq 1/2\}.$$

We define  $\varrho := e^{2\pi i/3}$ , so that  $-\bar{\varrho} = \varrho^2 = e^{4\pi i/3}$ . Pictorially, we have Figure 1

**2.1 PROPOSITION.** *The so defined  $D$  is a fundamental domain for the action of  $\Gamma$  on  $\mathbb{H}$ ; in particular we have that:*

1. every point in  $\mathbb{H}$  has a conjugate point in  $D$  with respect to the action;
2. if  $\tau, \tau' \in D$  are conjugate and different, then  $\Re\tau = \pm 1/2$  and  $\tau' = \tau \pm 1$ , or  $|\tau| = 1$ ,  $\tau' = -1/\tau$ ;
3. let  $\tau \in D$  and  $I(\tau) := |\{A \in \Gamma \mid A\tau = \tau\}|$ ; then  $I(\tau) = 1$  unless  $\tau = \iota$  ( $I(\iota) = 2$ ) or  $\tau \in \{\varrho, -\bar{\varrho}\}$  ( $I(\varrho) = I(-\bar{\varrho}) = 3$ ).

We can translate this situation into the stack language; the closure of  $\mathbb{H}/\Gamma$  can be identified with the weighted projective space  $\mathbb{P}(4, 6)$ ; so we can define a generic  $1/2 : 1$  map to  $\mathbb{P}^1$  (with two special points, corresponding to  $\iota$  and  $\varrho$ ).

**2.2 DEFINITION.** Let  $k \in \mathbb{Z}$ ; a weakly modular form of weight  $k$  is a meromorphic function  $f: \mathbb{H} \rightarrow \mathbb{C}$  such that  $f(A\tau) = (c\tau + d)^k f(\tau)$  for every  $A \in \text{SL}(2, \mathbb{Z})$ .

Let  $L$  be the space of meromorphic function  $f: \mathbb{H} \rightarrow \mathbb{C}$ ;  $\text{SL}(2, \mathbb{Z})$  acts on  $L$  in this way:

TODO

**2.3 COROLLARY.** *A function  $f$  is weakly modular of weight  $k$  if and only if  $f(-\tau^{-1}) = f(\tau)$  and  $f(\tau + 1) = f(\tau)$  (e.g. if it is invariant with respect to  $S$  and  $T$ ).*

Given  $\tau$ , write  $q := e^{2\pi i\tau}$ ; consider a weakly modular form of weight  $k$ ; then we can define  $\tilde{f}: E^* \rightarrow \mathbb{C}$  (where  $E^*$  is the punctured unit disk) by  $\tilde{f}(q) := f(\tau)$ . In particular,  $q \rightarrow 0$  corresponds to  $\tau \rightarrow \infty$ .

**2.4 DEFINITION.** We say that  $f$  is holomorphic at  $\infty$  if  $\tilde{f}$  is holomorphic at 0.

### 3. EXAMPLES

---

In other words,  $f$  is holomorphic at  $\infty$  if we can write  $\tilde{f} = \sum_{n \geq 0} a_n q^n$ , or, equivalently,  $f(\tau) = \sum_{n \geq 0} a_n (e^{2\pi i \tau})^n$ . This is how we make formal the request that  $f$  does not grow too fast for  $\tau \rightarrow \infty$ .

**2.5 DEFINITION.** Let  $k \in \mathbb{Z}$ ; a weakly modular form  $f$  of weight  $k$  is a *modular form of weight  $k$*  if  $f$  is holomorphic on  $\mathbb{H}$  and at  $\infty$ . In this case, we define  $f(\infty) := a_0$ ;  $f$  is called a *cuspidal form* if  $f(\infty) = 0$ .

Most interesting properties of modular forms are encoded in the Fourier coefficients  $a_n$ .

**2.6 REMARK.** Since  $-I \in \text{SL}(2, \mathbb{Z})$ , for a modular forms we have  $f(\tau) = f((-I)\tau) = (-1)^k f(\tau)$ ; in particular modular forms can exist only for  $k$  even.

## 3 EXAMPLES

**3.1 EXAMPLE (Eisenstein series).** Let  $\Lambda$  be a lattice in  $\mathbb{C}$ ; then  $\sum_{\lambda \in \Lambda} 1/|\lambda|^\sigma$  is convergent for every  $\sigma > 2$ . Let  $k \geq 2$ ; we define the Eisenstein series of weight  $k$  as

$$G_k(\tau) := \frac{(k-1)!}{2(2\pi i)^k} \sum'_{m,n \in \mathbb{Z}} \frac{1}{(m\tau + n)^k},$$

where the prime means that we exclude the value  $(0,0)$ . If  $k > 2$  the series is absolutely convergent, in the case  $k = 2$  we have to prove convergence with some other method. Assume  $k > 2$ ; then we can rearrange the terms of the series and this allow us to prove that  $G_k$  is a modular form of weight  $k$ . Indeed,

$$(c\tau + d)^k G_k(A\tau) = G_k(\tau)$$

since, disregarding multiplicative coefficients, we have

$$\begin{aligned} (c\tau + d)^k \sum_{m,n \in \mathbb{Z}} \frac{1}{(m(A\tau) + n)^k} &= \sum_{n,m \in \mathbb{Z}} \frac{1}{(m(a\tau + b) + n(c\tau + d))^k} = \\ &= \sum_{n,m \in \mathbb{Z}} \frac{1}{(m\tau + n)^k} \end{aligned}$$

since  $(a\tau + b, c\tau + d)$  is another basis of the lattice  $\mathbb{Z}\tau \oplus \mathbb{Z}$ . This tell us that  $G_k$  is a weakly modular form of weight  $k$  and holomorphic on  $\mathbb{H}$ ; the last thing to check is holomorphicity at  $\infty$ . Thanks to Euler identity, for  $z \in \mathbb{H}$  we have

$$\sum_{n \in \mathbb{Z}} \frac{1}{n+z} = \frac{\pi}{\tan(\pi z)} = -\pi i - 2\pi i \frac{e^{2\pi i z}}{1 - e^{2\pi i z}} = -\pi i - 2\pi i \sum_{n \geq 1} e^{2\pi i n z}.$$

We can also consider (it is no more than computing a derivative)

$$\sum_{n \in \mathbb{Z}} \frac{1}{(n+z)^k} = \frac{(-2\pi i)^k}{(k-1)!} \sum_{n \geq 1} n^{k-1} e^{2\pi i n z}.$$

Substituting in the Eisenstein series we get

$$\begin{aligned} G_k(\tau) &= \frac{(k-1)!}{2(2\pi i)^k} \left( \sum_{n \in \mathbb{Z}} \frac{1}{n^k} + \sum'_{m \in \mathbb{Z}} \sum_{n \in \mathbb{Z}} \frac{1}{(m\tau + n)^k} \right) = \\ &= \frac{(k-1)!}{(2\pi i)^k} \left( \sum_{n \geq 1} \frac{1}{n^k} + \sum_{m \geq 1} \sum_{n \in \mathbb{Z}} \frac{1}{(m\tau + n)^k} \right) = \\ &= \frac{(k-1)!}{(2\pi i)^k} \left( \sum_{n \geq 1} \frac{1}{n^k} + \sum_{m \geq 1} \frac{(-2\pi i)^k}{(k-1)!} \sum_{n \geq 1} n^{k-1} e^{2\pi i m n \tau} \right) = \\ &= \frac{(k-1)!}{(2\pi i)^k} \zeta(k) + \sum_{m \geq 1} \left( \sum_{d|m} d^{k-1} \right) q^m = \\ &= -\frac{B_k}{2k} + \sum_{n \geq 1} \sigma_{k-1}(n) q^n, \end{aligned}$$

where  $B_k$  is the  $k$ -th Bernoulli number and  $\sigma$  is the sum of divisor function; hence  $G_k$  is holomorphic at  $\infty$ . In particular

TODO

$$\begin{aligned} G_2(\tau) &= -\frac{1}{24} + q + 3q^2 + \dots, \\ G_4(\tau) &= \frac{1}{240} + q + 9q^2 + \dots \end{aligned}$$

For  $k = 2$ , the sum do not converge absolutely; we define

$$G_k^*(\tau) := -\frac{1}{8\pi} \lim_{\varepsilon \rightarrow 0} \sum_{n, m \in \mathbb{Z}} \frac{1}{(m\tau + n)^k |m\tau + n|^\varepsilon} = G_k(\tau) + \frac{1}{8\pi \Im \tau}.$$

The new series are absolutely convergent; but  $G_k^*$  is no more holomorphic since it depends explicitly on the imaginary part of  $\tau$ . We can compute how the transformation property behaves on the correction term:

$$G_k(A\tau) = (c\tau + d)^k G_k(\tau) - \frac{c(c\tau + d)}{4\pi i}.$$

**3.2 EXAMPLE (Discriminant function).** We can define  $\Delta: \mathbb{H} \rightarrow \mathbb{C}$ , the *discriminant function*, as  $\Delta(\tau) := q \prod_{j \geq 1} (1 - q^j)^{24}$ , where as usual  $q = e^{2\pi i \tau}$ . This converges on  $\mathbb{H}$ ; if it is modular, then it is a cusp form. Obviously  $\Delta(\tau + 1) = \Delta(\tau)$ ; define  $\Delta' := \frac{\partial \Delta}{\partial \tau}$  and let  $\Delta'/\Delta(\tau)$  the logarithmic derivative of  $\Delta$ . We find

that

$$\frac{\Delta'}{\Delta}(\tau) = 2\pi i \left( 1 - 24 \underbrace{\sum_{n \geq 1} \frac{nq^n}{1-q^n}}_{\sum_{n \geq 1} \sum_{i \geq 1} iq^{ni} = \sum_{m \geq 1} \sigma_k(m)q^m} \right) = -24 \cdot 2\pi i G_k(\tau).$$

Then

$$\frac{d}{d\tau} \log \left( \Delta \left( -\frac{1}{\tau} \right) \right) = \frac{1}{\tau^2} \frac{\Delta'}{\Delta} \left( -\frac{1}{\tau} \right) = \frac{\Delta'}{\Delta}(\tau) + \frac{12}{\tau} = \frac{d}{d\tau} \log(\Delta(\tau)\tau^{12}),$$

that is  $\Delta(-\tau^{-1}) = \text{const} \cdot \tau^{12} \Delta(\tau)$ . If we put  $\tau = i$ , then  $-\tau^{-1} = i$  and  $\tau^{12} = 1$ , so the constant must be 1 and  $\Delta$  is a cusp form of weight 12.

Lecture 2 (2 hours)  
March 12<sup>th</sup>, 2009

3.3 REMARK. We denote the vector space of modular forms of weight  $k$  with  $M_k$  and the vector space of cusp forms of weight  $k$  with  $S_k$ . It is obvious that if  $f_k \in M_k$  and  $f_l \in M_l$  then  $f_k f_l \in M_{k+l}$ .

#### 4 ZEROES OF MODULAR FORMS

If  $f$  is a meromorphic function on  $\mathbb{H}$  we can define its order at a point  $p \in \mathbb{H}$  as  $v_p(f)$ , the integer such that  $f(\tau)(\tau - p)^{-v_p(f)}$  is holomorphic and non-zero at  $p$ . If  $f$  is a modular form, then  $f(\tau) = (c\tau + d)^{-k} f(A\tau)$  for every  $A \in \text{SL}(2, \mathbb{Z})$ , so  $v_p(f) = v_{Ap}(f)$  for every  $A \in \text{SL}(2, \mathbb{Z})$ . In particular, if  $f = \sum_{n \geq 0} a_n q^n$ , then we define  $v_\infty(f) := v_0(\tilde{f})$ .

4.1 THEOREM. Let  $f$  be a modular form of weight  $k$ , then

$$(3) \quad v_\infty(f) + \frac{1}{2}v_i(f) + \frac{1}{3}v_\rho(f) + \sum_{p \in \mathbb{H}/\Gamma \setminus \{0, i\}} v_p(f) = \frac{k}{12}.$$

In the stack interpretation, we define modular forms as sections of a line bundle  $\mathcal{L}_2 \rightarrow \overline{M}_{1,1} \cong \mathbb{P}(4, 6)$ ; then the theorem says that the degree of this line bundle is  $k/12$ .

*Proof.* We can assume that our modular form  $f$  has no zeroes on the boundary of the fundamental domain  $D$  (except maybe in  $i$  or  $\rho$ ), since we can move slightly  $D$  until this is true.

Now we can integrate  $df/f$  on the boundary of  $D$ . More formally, consider Figure 2a: first, we integrate on a path like  $\gamma$  in such a way that all internal singularities are inside  $\gamma$ ; by the residue theorem,

$$\frac{1}{2\pi i} \int_\gamma \frac{df}{f} = \sum_{p \in \mathbb{H}/\Gamma \setminus \{0, i\}} v_p(f).$$

We will compute now the same integrals piece by piece. For simplicity, we forget about the coefficient  $2\pi i$ .




Figure 2: Proof of Theorem 4.1.

- The integral on the arc near  $q$  is just  $-1/6v_q(f)$ , since we can compute the integral along the path  $\gamma_q$  of Figure 2b getting  $-v_q(f)$  (since we are going clockwise this time) and then, passing to the limit of the radius, we have to divide by 6 since the angle is  $\pi/3$ .
- The same applies to the integral on the arc near  $q^2$ .
- With the same method, the integral on the arc near  $\iota$  is  $-1/2v_\iota(f)$ .
- Using the transformation  $\tau \mapsto q$ , the horizontal segment becomes a whole clockwise circle around  $q = 0$ , so the integral on the segment is  $-v_\infty(f)$ .
- The two vertical path are obtained one from the other by applying  $T$  or  $T^{-1}$ ; since  $f(T\tau) = f(\tau)$  and they are in opposite direction, the sum of the two integrals is 0.
- The two remaining arcs are obtained one from the other by applying  $S$  or  $S^{-1}$ ; this time,  $f(S\tau) = \tau^k f(\tau)$ , so

$$\frac{df(S\tau)}{f(S\tau)} = k \frac{d\tau}{\tau} + \frac{df(\tau)}{f(\tau)};$$

then, the sum of the two integral is

$$\int \left( \frac{df(\tau)}{f(\tau)} - \frac{df(S\tau)}{f(S\tau)} \right) = \int -k \frac{dz}{z} = -k \left( -\frac{1}{12} \right) = \frac{k}{12}.$$

Comparing the two results we get

$$\sum_{p \in \mathbb{H}/\Gamma \setminus \{q, \iota\}} v_p(f) = -\frac{1}{3}v_q(f) - \frac{1}{2}v_\iota(f) - v_\infty(f) + \frac{k}{12}. \quad \square$$

We recall that  $M_k = 0$  for  $k$  odd, that is, there are no odd weighted modular forms; moreover, since  $G_{2k} \in M_{2k}$  is a modular forms that is not a cusp form

(Bernoulli numbers are always non-zero) it follows that  $\dim M_{2k}/S_{2k} \geq 1$ ; but  $S_{2k}$  is the kernel of the map  $f \mapsto f(\infty)$ , so  $\dim M_{2k}/S_{2k} \leq 1$ ; hence,  $M_{2k} = S_{2k} \oplus \mathbb{C}G_{2k}$ .

4.2 THEOREM.

1. If  $k < 0$  or  $k$  is odd, then  $M_k = 0$ .
2. For  $k \in \{0, 4, 6, 8, 10\}$ ,  $S_k = 0$  and  $M_k = \mathbb{C}G_k$ ;  $M_2 = 0$ ;  $G_0 = 1$ .
3. Multiplication by  $\Delta$  gives an isomorphism  $M_{k-12} \rightarrow S_k$  for all  $k$ .

*Proof.* The first statement follows from equation (3), since all left-hand side terms are non-negative. We have  $M_2 = 0$  since  $1/6$  cannot be written as a non-negative integral combination of  $1, 1/2$  and  $1/3$ ;  $S_k = 0$  for  $k < 12$  is trivial since for a cusp form we have  $v_\infty(f) \geq 1$ .

Since  $\Delta$  has no zeroes on  $\mathbb{H}$ , if  $f \in S_k$  we can write  $g := f/\Delta$  and  $g$  has weight  $k - 12$ . Now  $v_p(g) = v_p(f)$  for every  $p \in \mathbb{H}$  and  $v_\infty(g) = v_\infty(f) - 1$ , hence  $g \in M_{k-12}$ . From this it follows the rest of the second statement.  $\square$

4.3 COROLLARY. The dimension of  $M_k$  is

$$\dim M_k = \begin{cases} 0 & \text{if } k < 0 \text{ or } k \text{ odd;} \\ \lfloor k/12 \rfloor & \text{if } k \equiv 2 \pmod{12}; \\ \lfloor k/12 \rfloor + 1 & \text{if } k \not\equiv 2 \pmod{12}. \end{cases} \quad (12).$$

4.4 COROLLARY. Let  $M_A := \bigoplus_k M_k$ ; then as a graded ring  $M_A \cong \mathbb{C}[G_4, G_6]$ . Equivalently, a basis of  $M_k$  is  $\{G_4^a G_6^b \mid 4a + 6b = k\}$ .

*Proof.* In multiple steps.

- If  $k \leq 6$  this is obvious.
- Since  $M_{12} = \mathbb{C}G_{12} \oplus \Delta$ , and we have  $\lambda_4 G_4 + \lambda_6 G_6 \in M_{12}$  for every  $\lambda_4, \lambda_6 \in \mathbb{C}$ , then the statement is true for  $M_{12}$  and in particular  $\Delta$  is generated by  $G_4$  and  $G_6$ .
- By induction on even  $k$  greater than 6: choose  $a$  and  $b$  such that  $4a + 6b = k$  and let  $g := G_4^a G_6^b \in M_k$ ;  $g$  is not a cusp form, so for every  $f \in M_k$  there exists  $\lambda \in \mathbb{C}$  such that  $f - \lambda g$  is a cusp form; but then  $f - \lambda g \in S_k = M_{k-12}\Delta$  and we conclude since both  $\Delta$  (by the previous point) and  $M_{k-12}$  (by induction) are generated by  $G_4$  and  $G_6$ .  $\square$

Define now  $E_k := G_k \cdot (-2k/B_k) = 1 + \dots$ .

4.5 COROLLARY.

$$E_4^2 = E_8.$$



---

By this corollary we can state the following non-trivial identity for every  $n > 0$ :

$$\sigma_7(n) = \sigma_3(n) + 120 \sum_{m=1}^{n-1} \sigma_3(m)\sigma_3(n-m).$$

Another identity is  $E_4^3 - E_6^2 = 1728\Delta$ .

## 5 THETA FUNCTIONS

Let  $\Lambda$  be a lattice in  $\mathbb{R}^n$ , such that  $v \cdot v \in \mathbb{N}$  for every  $v \in \Lambda$ . We wonder how many vectors of a given length exist in  $\Lambda$ . We define a generating function

$$\Theta_\Lambda(\tau) = \sum_{n \geq 0} |\{v \in \Lambda \mid v \cdot v = n\}| q^{n/2},$$

where again  $q = e^{2\pi i \tau}$ . We can write the same function in a simpler way:  $\Theta_\Lambda(\tau) = \sum_{v \in \Lambda} q^{v \cdot v/2}$ . We want to show that these are modular forms; to do this we make use of the Poisson summation formula.

Let  $\varphi: \mathbb{R}^n \rightarrow \mathbb{R}$  a smooth function rapidly decreasing at  $\infty$ , that is, such that as  $\|x\| \rightarrow \infty$ , it goes as  $\|x\|^{-c}$  for  $c \geq n$ . The Fourier transform of  $\varphi$  is  $\hat{\varphi}: \mathbb{R}^n \rightarrow \mathbb{R}$  defined by

$$\hat{\varphi}(t) = \int_{\mathbb{R}^n} \varphi(x) e^{-2\pi i t x} dx.$$

Let  $\mu$  the volume of  $\mathbb{R}^n/\Lambda$  (equivalent to  $\det(a_i \cdot a_j)^{n/2}$  where  $a_i$  is a basis of  $\Lambda$ ); let  $\Lambda^\vee$  be the dual lattice, that is the set of all  $w \in \mathbb{R}^n$  such that  $w \cdot v \in \mathbb{Z}$  for every  $v \in \Lambda$ .

**5.1 THEOREM** (Poisson summation formula).

$$\sum_{v \in \Lambda} \varphi(v) = \frac{1}{\mu} \sum_{w \in \Lambda^\vee} \hat{\varphi}(w).$$

Let  $t \in \mathbb{R}_{>0}$  and define  $\tilde{\Theta}_\Lambda(t) := \sum_{v \in \Lambda} e^{-\pi t v \cdot v}$ .

**5.2 PROPOSITION.**

$$\tilde{\Theta}_{\Lambda^\vee}(t^{-1}) = t^{n/2} \mu \tilde{\Theta}_\Lambda(t).$$

*Proof.* Fix  $t$  and put  $f(x_1, \dots, x_n) := e^{-\pi(x_1^2 + \dots + x_n^2)}$ . It is easy to prove that  $f$  is a rapidly decreasing function and that  $\tilde{f} = f$ . Consider the lattice  $\sqrt{t}\Lambda$ ; its dual is  $1/\sqrt{t}\Lambda^\vee$  and its volume is  $t^{n/2}\mu$ .

Applying the Poisson summation formula, we get

$$\sum_{v \in \Lambda} e^{-\pi t v \cdot v} = \frac{t^{-n/2}}{\mu} \sum_{w \in \Lambda^\vee} e^{-\pi^2 t w \cdot w}.$$

This gives the statement. □

Assume from now on that  $\Lambda$  is a unimodular, even, integral lattice, that is, such that  $\Lambda^\vee = \Lambda$ ,  $v \cdot v \in 2\mathbb{Z}$  and  $w \cdot v \in \mathbb{Z}$  for every  $v, w \in \Lambda$ .

5.3 THEOREM.

1.  $\Theta_\Lambda(\tau) = \sum_{v \in \Lambda} q^{v \cdot v/2}$  is a modular form of weight  $n/2$ ;
2.  $n$  is divisible by 8.

*Proof.* Since  $v \cdot v \in 2\mathbb{Z}$ , the definition of  $\Theta_\Lambda(\tau)$  is a  $q$ -development; moreover it is clear that it is invariant under  $\tau \rightarrow \tau + 1$ . We want to prove that  $\Theta_\Lambda(-1/\tau) = (\iota\tau)^{n/2} \Theta_\Lambda(\tau)$ ; this is enough because, if  $8 \mid n$ , the  $\iota$  go away and we remain with a modular form. Since  $\Theta_\Lambda$  is an analytic function, we can prove it just for  $\tau = \iota t$  with  $t \in \mathbb{R}_{>0}$ . Now,  $\Theta_\Lambda(\iota t) = \sum_{v \in \Lambda} e^{-\pi t v \cdot v} = \tilde{\Theta}_\Lambda(t)$ ; besides,  $\Theta_\Lambda(-1/\iota t) = \tilde{\Theta}_\Lambda(-1/t)$ . The statement then follows from Proposition 5.2.

Conversely, assume  $8 \nmid n$ ; replacing  $\Lambda$  by  $\Lambda^2$  or  $\Lambda^4$  we may assume that  $n \equiv 4 \pmod{8}$ , so  $\Theta_\Lambda(-1/\tau) = -\tau^{n/2} \Theta_\Lambda(\tau)$ . We recall that from every function  $f$  on  $\mathbb{H}$  we can define  $f|_{k,A}(\tau) = (c\tau + d)^{-k} f(A\tau)$  for  $A \in \text{SL}(2, \mathbb{Z})$ . In particular, we apply this to  $f = \Theta_\Lambda$ ,  $k = n/2$  and  $A \in \{S, T\}$ . We obtain respectively  $-\Theta_\Lambda(\tau)$  and  $\Theta_\Lambda(\tau)$ ; but  $(ST)^3 = I$ , so

$$\Theta_\Lambda(\tau) = \Theta_\Lambda|_{n/2, (ST)^3} = -\Theta_\Lambda(\tau),$$

TODO

contraddiction. □

5.4 COROLLARY. *There is a cusp form  $f_\Lambda$  of weight  $n/2$  such that  $\Theta_\Lambda = E_{n/2} + f_\Lambda$ .*

For  $n \equiv 0 \pmod{8}$  it is quite easy to define a unimodular, even, integral lattice on  $\mathbb{R}^n$ . For example, start with the lattice  $B_n := \{v \in \mathbb{Z}^n \mid v \cdot v \in 2\mathbb{Z}\}$  and consider  $\Lambda_n := B_n \oplus (1/2, \dots, 1/2)\mathbb{Z}$ . This construction gives in particular  $\Lambda_8 = E_8$ .

5.5 EXAMPLE. We have  $\Theta_{\Lambda_8} = E_4$ , since there is no cusp forms of weight 4. Besides,  $E_4 = 1 + 240 \sum_{n \geq 1} \sigma_3(n) q^n$  and this gives us the number of lattice with the properties we wanted. In the same way,  $\Theta_{\Lambda_{16}} = \Theta_{\Lambda_8 \oplus \Lambda_8} = E_4^2 = E_8$ .

## 6 MODULAR FORMS FOR CONGRUENCE SUBGROUPS

The group  $\text{SL}(2, \mathbb{Z})$  contains copies of the integers: they are identified with the subgroups  $\Gamma(N)$  of matrices  $A \equiv I \pmod{N}$ ; we have also the subgroups

$$\begin{aligned} \Gamma^0(N) &:= \{A \in \text{SL}(2, \mathbb{Z}) \mid A \equiv \begin{pmatrix} * & 0 \\ * & * \end{pmatrix} \pmod{N}\}, \\ \Gamma_0(N) &:= \{A \in \text{SL}(2, \mathbb{Z}) \mid A \equiv \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N}\}. \end{aligned}$$

6.1 DEFINITION. A subgroup  $G$  of  $\text{SL}(2, \mathbb{Z})$  is called a *congruence subgroup* if  $\Gamma(N) \subseteq G$

---

6.2 DEFINITION. Fixed a congruence subgroup  $G$ , a holomorphic function  $f: \mathbb{H} \rightarrow \mathbb{C}$  is called a *modular form of weight  $k$  on  $G$*  if:

1.  $f|_{k,A} = f$  for every  $A \in G$  (that is,  $f(\tau) = (c\tau + d)^{-k} f(A\tau)$ );
2.  $f$  is holomorphic at the cusps: for every  $A \in \text{SL}(2, \mathbb{Z})$ , there exists  $l > 0$  such that  $f|_{k,A} = \sum_{n \geq 0} a_n q^{n/l}$  with  $a_n \in \mathbb{C}$  and  $q^{n/l} = e^{2\pi i \tau n/l}$ .

There is a geometric interpretation of the second condition.

- Let  $\bar{\mathbb{Q}} := \mathbb{Q} \cup \{\infty\}$ ; the action of  $\text{SL}(2, \mathbb{Z})$  on  $\mathbb{H}$  extends to  $\bar{\mathbb{Q}}$  by  $A\alpha := \frac{a\alpha+b}{c\alpha+d}$  (these action sends  $\bar{\mathbb{Q}}$  to itself). A cusp of  $\mathbb{H}/G$  is an element of  $\bar{\mathbb{Q}}/G$ ; in particular, if  $G = \text{SL}(2, \mathbb{Z})$  we have only one cusp which we can imagine to be  $\infty$ . In general,  $\mathbb{H}/G$  can be compactified to a complete orbifold Riemann surface as  $\bar{\mathbb{H}}/G = \mathbb{H}/G \cup \{\text{cusps}\}$ .
- Let  $\alpha \in \bar{\mathbb{Q}}$  and  $A \in \text{SL}(2, \mathbb{Z})$ , with  $A(\infty) = \alpha$ . Let  $l \geq 0$  such that  $T^l \in A^{-1}GA$ ; then

$$(f|_{k,A})|_{k,T^l} = f|_{k,AT^l} = f|_{k,A},$$

that is,  $f|_{k,A}$  is mapped to itself by  $\tau \rightarrow \tau + l$ . We fix  $l$  to be minimal with respect to his condition; this  $l$  is called *width* of the cusp. Now we can write  $f|_{k,A} = \sum_{n \in \mathbb{Z}} a_n q^{n/l}$ , and holomorphic at cusp  $\alpha$  is equivalent to  $a_n = 0$  for every  $n < 0$ .

- Geometrically,  $\bar{\mathbb{H}}/G$  is a complex orbifold that has an obvious map  $\varphi$  to  $\bar{\mathbb{H}}/\text{SL}(2, \mathbb{Z})$ ; this map is a branch cover of degree  $[\text{SL}(2, \mathbb{Z}) : G]$ . The point  $\infty$  in the target has as fiber the set of cusps in the source; moreover, the order of  $\varphi$  at a cusp is just its width (that is, at  $\alpha$ ,  $q^{n/l}$  is a local coordinate).

6.3 EXAMPLE. Consider  $G := \Gamma(2)/\{\pm I\}$ ; it can be proved that it is the free group  $\langle (\frac{1}{0} \frac{2}{1}), (\frac{1}{2} \frac{0}{1}) \rangle$ . A fundamental domain is represented in Figure 3. Its cusps then are  $0, 1, \infty$ ; the width of  $\infty$  is 2. As before, we define the set of modular forms to be  $M_{k,G}$  with the subspace  $S_{k,G}$  of cusp forms (that is, modular forms such that  $f(\alpha) = 0$  for every cusp  $\alpha$ ). They are finite dimensional vector spaces and we can compute their dimensions.

6.4 EXAMPLE. The theta function  $\Theta_{\mathbb{Z}^4}$  is  $\sum_{n_1, \dots, n_4 \in \mathbb{Z}} q^{\sum n_i^2}$ . This is not even, so it is not a modular form; but a similar argument of the one did before shows that it is a modular form on some subgroup, precisely a modular form of weight 2 on  $\Gamma^0(4)$ .

6.5 COROLLARY.

1. Every positive integer is the sum of four squares;
2.  $|\{n_1, \dots, n_4 \in \mathbb{Z} \mid \sum n_i^2 = n\}| = 8(\sum_{d|n, 4 \nmid d} d)$ .




Figure 3: A fundamental domain of the action of  $G$  on  $\mathbb{H}$ .

*Proof.* The first statement is obvious; for the second, consider  $8G_2(\tau) - 32G_2(4\tau)$ ; this is a modular form of weight 2 on  $\Gamma^0(4)$ . This is quite surprising since  $G_2$  is not even; but we recall that  $G_2^*(\tau)$  is not holomorphic but transforms as a modular forms; so the one we are considering is just  $8G_2^*(\tau) - 32G_2^*(4\tau)$ .  $\square$

## 7 HECKE THEORY

On modular forms there is an algebra of operators (the Hecke operators) such that there is a basis of simultaneous eigenvalues for the operators.

Recall that we have an isomorphism of vector spaces between:

- complex functions  $F$  of oriented lattices  $\Lambda = \mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2$  with  $\Im(\omega_1/\omega_2) > 0$  such that  $F(a\Lambda) = a^{-k}\Lambda$ ;
- holomorphic functions  $f: \mathbb{H} \rightarrow \mathbb{C}$  such that  $f(A\tau) = (c\tau + d)^{-k}f(\tau)$  for every  $A \in \mathrm{SL}(2, \mathbb{Z})$ .

In particular we associate to a morphism  $F$  the function  $f(\tau) := F(\tau\mathbb{Z} \oplus \mathbb{Z})$  and to a function  $f$  the morphism such that  $F(\mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2) := \omega_2^{-k}f(\omega_1/\omega_2)$ .

Let  $F$  be a lattice function of weight  $k$ ; define the operators  $T_n = T_{n,k}$  by

$$T_n F(\Lambda) := n^{k-1} \sum_{\Lambda' \subseteq \Lambda, [\Lambda:\Lambda'] = n} F(\Lambda').$$

These  $T_n$  have an interpretation as morphisms of moduli space of elliptic curves with additional level structure. Note anyway that  $T_n F$  is a lattice function of weight  $k$ . Thus, denoting the corresponding function with  $f: \mathbb{H} \rightarrow \mathbb{C}$ , we define  $T_n f(\tau) := T_n F(\tau\mathbb{Z} \oplus \mathbb{Z})$ . Then for  $T_n F$  to be a lattice function of weight  $k$  means that  $T_n f(A\tau) = (c\tau + d)^{-k} T_n f$ . After some computation we obtain a description in terms of  $\tau$ :  $T_n f(\tau) = n^{k-1} \sum_{A \in \Gamma \backslash \mathcal{M}_n} (c\tau + d)^{-k} f(A\tau)$ . The summation indices means that  $A$  runs through a system of representatives of  $\Gamma \backslash \mathcal{M}_n$ , where  $\mathcal{M}_n$  is the set of  $2 \times 2$  matrices with entries in  $\mathbb{Z}$  and determinant  $n$ , and  $\mathrm{SL}(2, \mathbb{Z})$  acts on  $\mathcal{M}_n$  by multiplication on the left.

If  $f \in M_k$ , then  $T_n f$  is holomorphic on  $\mathbb{H}$ ; plus, we already seen that it transforms as a modular forms; to check that  $T_n f$  is a modular form, we

need to prove that it is holomorphic at  $\infty$ ; we do this writing down its  $q$ -development.

7.1 THEOREM.

1. Let  $f \in M_k$  with Fourier development  $f(\tau) = \sum_{n \geq 0} c(n)q^n$ ; then

$$T_n f(\tau) = \sum_{m \geq 0} \left( \sum_{d|n, d|m, d \geq 0} d^{k-1} \right) c(nm/d^2) q^m.$$

In particular,  $T_n f \in M_k$  and if  $f$  is a cusp form, then  $T_n f$  is.

2.  $T_n$  satisfies

$$T_n T_m = \sum_{d|n, d|m, d \geq 0} d^{k-1} T_{nm/d^2};$$

in particular,  $T_n$  and  $T_m$  commutes and if  $(m, n) = 1$ ,  $T_n T_m = T_{nm}$ .

*Proof.* A system of representatives of  $\Gamma \backslash \mathcal{M}_n$  is the set of matrices  $\begin{pmatrix} a & b \\ 0 & d \end{pmatrix}$  such that  $ad = n$  and  $0 \leq b < d$ . Then  $T_n f(\tau) = n^{k-1} \sum_{a, d > 0, ad=n} \sum_{b=0}^{d-1} d^{-k} f\left(\frac{a\tau+b}{d}\right)$ . Substituting the  $q$ -development of  $f$  we obtain

$$T_n f(\tau) = n^{k-1} \sum_{a, d > 0, ad=n} \sum_{b=0}^{d-1} d^{-k} \sum_{m \geq 0} c(m) e^{2\pi i m \frac{a\tau+b}{d}}.$$

Note that

$$\sum_{b=0}^{d-1} e^{2\pi i mb/d} = \begin{cases} 0 & d \nmid m \\ d & d \mid m \end{cases}$$

Now, observe that the second statement follows from the first by easy computations.  $\square$

7.2 REMARK. Observe that  $T_p T_{p^n} = T_{p^{n+1}} + p^{k-1} T_{p^{n-1}}$  when  $p$  is prime; then if  $n = p_1^{n_1} \cdots p_l^{n_l}$ , then  $T_n = T_{p_1^{n_1}} \cdots T_{p_l^{n_l}}$ .

7.3 DEFINITION. The Hecke operators are the operators  $T_n: M_k \rightarrow M_k$ .

The Hecke operators are a set of commuting linear map; it is possible then to search for common eigenvectors, that is, modular forms  $f$  such that  $T_n f = \lambda_n f$  for every  $n \geq 1$ .

7.4 DEFINITION. Let  $f \in M_k$  be a common eigenvector of all  $T_n$ ; assume  $f(\tau) = \sum_{n \geq 0} a_n q^n$  with  $a_n = 1$ ; then  $f$  is called a Hecke form.

The condition on  $a_n = 1$  is to normalize the form. At first it appears that the parameter  $q$  is not so special; but it happens that the coefficients of the Fourier transform have a geometrical meaning; in particular, they are related to the eigenvalues.

## 8. L-SERIES

---

7.5 COROLLARY. Let  $f = \sum_{n \geq 0} a_n q^n$  be a Hecke form; then  $T_n f = a_n f$  for all  $n \geq 1$ .

7.6 COROLLARY. Let  $f = \sum_{n \geq 0} a_n q^n$  be a Hecke form; then  $a_n a_m = \sum_{d|n, d|m} d^{k-1} c(nm/d^2)$ . In particular, if  $(n, m) = 1$ , then  $a_n a_m = a_{nm}$ .

*Proof.* It is obvious by the formula of  $T_n f(\tau)$ . Since  $T_n f(\tau) = \lambda_n f(\tau)$  and  $a_1 = 1$ , we can write

$$\lambda_n = \lambda_n a_1 = \sum_{d|n, d|1} d^{k-1} c(nm/d^2) = a_n. \quad \square$$

7.7 EXAMPLE. The Eisenstein series  $G_k$  is a Hecke form for  $k \geq 4$ ; so it is  $\Delta$ .

7.8 COROLLARY.

$$\tau(n)\tau(m) = \sum_{d|n, d|m} d^{11} \tau(nm/d^2).$$

7.9 THEOREM. The Hecke forms form a basis for  $M_k$  for all  $k$ .

## 8 L-SERIES

Let  $f := \sum_{n \geq 0} a_n q^n$  be a Hecke form; we can associate to it its *L-series*

$$L(f, s) := \sum_{n \geq 1} \frac{a_n}{n^s}$$

that converges absolutely and uniformly for  $\Re s > k$  and is a holomorphic function for  $\Re s > 0$ . To prove these convergency results we need some machinery.

8.1 LEMMA. Let  $f := \sum_{n \geq 0} a_n q^n \in M_k$ ; then  $a_n \in O(n^{k-1})$  (that is,  $a_n/n^{k-1} \rightarrow 0$  as  $n \rightarrow \infty$ ).

8.2 COROLLARY. We can write  $L(f, s)$  as an Euler product:

$$L(f, s) = \prod_{p \text{ prime}} \frac{1}{1 - a_p p^{-s} + p^{k-1-2s}}.$$

*Proof.* From  $a_n a_m = a_{nm}$  if  $(n, m) = 1$ , it follows that  $a_{p_1^{n_1} \dots p_l^{n_l}} = a_{p_1^{n_1}} \dots a_{p_l^{n_l}}$ ; then  $L(f, s) = \prod_{p \text{ prime}} \sum_{n \geq 0} a_p^n p^{-ns}$ . We have to show that  $\sum_{n \geq 0} a_p^n p^{-ns} = (1 - a_p p^{-s} + p^{k-1-2s})^{-1}$ . We know that  $a_{p^{n+1}} - a_p a_{p^n} + p^{k-1} a_{p^{n-1}} = 0$  for  $p$  prime; if we multiply the series with  $1 - a_p p^{-s} + p^{k-1-2s}$ , we see that the constant term, with respect to  $t := p^{-s}$  is 1, that the second term is 0 and by induction we get that all other coefficients are 0 using the previous relation.  $\square$

8.3 EXAMPLE. We can write the Riemann Zeta function  $\zeta(s) = \sum_{n \geq 1} 1/n^2$  as  $\prod_{p \text{ prime}} 1/(1 - p^{-s})$ . We can compute  $L(G_k, s)$ ; if  $p$  is a prime,  $\sigma_{k-1}(p) = 1 + p^{k-1}$

---

and the denominator of the terms of the L-series is  $1 + \sigma_{k-1}(p)p^{-s} + p^{k-1-2s} = (1 - p^{k-1-s})(1 - p^{-s})$  and it follows that

$$L(G_k, s) = \prod_{p \text{ primes}} \left( \frac{1}{1 - p^s} \right) \left( \frac{1}{1 - p^{k-1-s}} \right) = \zeta(s)\zeta(s - k + 1).$$

**8.4 THEOREM.** *If  $f$  is a Hecke form of weight  $k$ , then  $L(f, s)$  has a meromorphic continuation to the whole  $\mathbb{C}$  and satisfies some functional equation; if  $f$  is a cusp form, then  $L(f, s)$  is an entire function; otherwise it has a simple pole at  $s = k$ .*

#### REFERENCES

[Ser73] J.-P. Serre, *A course in arithmetic*, Springer-Verlag, New York, 1973, Translated from the French, Graduate Texts in Mathematics, No. 7.